

Antioxidant, iron chelating, lipase inhibition activities and metabolite's prediction of hydroethanolic leaf extract of *Conocarpus erectus*

^{1,*}Raza, S.A., ¹Chaudhary, A.R., ²Mumtaz, M.W. and ³Bashir, S.

¹Department of Chemistry, Government College University, Lahore, Pakistan

²Department of Chemistry, University of Gujrat, Pakistan

³Division of Science and Technology, University of Education, College Road Lahore, Pakistan

Article history:

Received: 22 August 2019

Received in revised form: 16 September 2019

Accepted: 18 September 2019

Available Online: 14 October 2019

Keywords:

Antioxidant,
Fe-chelating,
Lipase inhibition,
Ultrasonication,
Conocarpus erectus,
¹H-NMR.

DOI:

[https://doi.org/10.26656/fr.2017.4\(2\).300](https://doi.org/10.26656/fr.2017.4(2).300)

Abstract

The current study was conducted to evaluate the antioxidant, Fe-chelating and lipase inhibition activities of hydroethanolic leaf extracts of *Conocarpus erectus*. The proton magnetic resonance spectroscopy (¹H-NMR) was used to identify the major types of primary and secondary metabolites in extract. The 60% ethanolic extract was found to be the most effective fraction regarding antioxidant, Fe chelating and lipase inhibitory properties. The 60% ethanolic leaf extract exhibited total antioxidant power of 223.0±3.27 mg ASE/g PE, β-carotene bleaching inhibition of 81.39±2.11%, Fe-chelating of 68.21±1.17% and pancreatic lipase inhibition of 50.60±1.47%, respectively. The ¹H-NMR-based prediction of metabolites provided the information for presence of polyphenolic secondary metabolites and organic acids which might be responsible for the biological activities of extract. These findings of this work may be extended for metabolite characterization, *in vivo* trials and toxicity determination to get benefits for functional food development with antioxidative and antiobesity attributes.

1. Introduction

Reactive oxygen species (ROS) are produced in the human body as result of metabolic functions and play an important role in cell signaling. The overproduction of ROS is counterbalanced by antioxidant defense system of body. Any imbalance in ROS and antioxidants may create a situation of oxidative stress which involves initiation and progression of many chronic health disorders. The antioxidant intake is considered an effective tool to mitigate and manage oxidative stress and stress-related disorders like diabetes, cancer, obesity, aging and many others (Piyush *et al.*, 2010; William *et al.*, 2018). The commercially available synthetic antioxidants are although very effective but pose serious threats to human health due to their toxic nature. Commonly used synthetic antioxidants like butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT) and ter-butyl hydroquinone (TBHQ) have been proved toxic (Madhujith and Shahidi, 2006; Wang *et al.*, 2010). There are still many concerns about the safety of synthetic antioxidants which are being monitored regularly in Europe and developed countries.

Under such circumstances, plants provide a wide range of opportunities for availability of potent

antioxidants of safe and therapeutic nature. The plant extracts rich in polyphenols were reported to have notable antioxidant and enzyme inhibition properties. These activities were indicators of possible medicinal attributes of plants (Raza *et al.*, 2013; William *et al.*, 2018). The iron-chelating activity of plant extracts is reported as efficient strategy to mitigate the harmful impacts of overloaded iron in body. The iron overload is involved in production of ROS, initiation and progression of diabetes along with damage to liver function (Shaaban *et al.*, 2016). Similarly, pancreatic lipase inhibition is a useful way to reduce the burden of obesity. Plants have been reported as a natural source for safe metal chelating and lipase inhibitory agents. The antioxidants present in plants were mentioned as potent metal chelators and natural inhibitors of pancreatic lipase.

Conocarpus erectus (Combretaceae) has gained importance due to recently explored medicinal aspects including antidiabetic, antimicrobial, anticancer and hepatoprotective effects (Saleh *et al.*, 2012; Saleh *et al.*, 2013; Nascimento *et al.*, 2016; Raza *et al.*, 2018). The metal chelating activity and pancreatic lipase inhibitory properties of *C. erectus* have not been reported previously as per best of available information.

*Corresponding author.

Email: chemstone@yahoo.com

Therefore, the current work was performed to evaluate the antioxidant activity, iron-chelating activity, pancreatic lipase inhibition potential of *C. erectus* leaf extracts along with ¹H-NMR based metabolite prediction.

2. Material and methods

2.1 Hydroethanolic extraction

The *C. erectus* leaves were identified from the Department of Botany, GCU Lahore vide voucher specimen GC.Herb.33.Bot. 3379. Fresh leaves were collected and immediately quenched with liquid nitrogen, lyophilized and suspended in solvent systems (aqueous, 20% ethanol, 40% ethanol, 60% ethanol and pure ethanol). The samples were ultrasonicated at Soniprep 150, filtered and extra solvent was removed on rotary evaporator under reduced pressure. The extracts were stored at low temperature until further use.

2.2 Total antioxidant power

Total antioxidant activity of all extract fractions was based on phosphomolybdenum complex formation methods with little modification as previously reported (Prieto *et al.*, 1999). Briefly, 250 µg/mL of each extract was mixed with 4 mL of reagent solution consisting of 0.6 M sulphuric acid, 4 mM ammonium molybdate and 28 mM sodium phosphate in capped plastic vials. The obtained mixtures along with a blank solution (4 ml reagent solution) were incubated in water bath at 95°C for 90 mins. After cooling to room temperature, absorbance was noted at 695 nm. A standard curve was drawn using ascorbic acid. The butylated hydroxyanisole (BHA) was used as standard antioxidant for comparison of antioxidant potential of extracts. The results were represented as ascorbic acid equivalent per gram dried plant extract (ASE/g DE).

2.3 β-carotene bleaching assay

Antioxidant activity of *C. erectus* leaf extracts was determined by assessing the bleaching of β-carotene (yellow color) in the presence of linoleic acid (Shon *et al.*, 2003). Initially, 2 mg of β-carotene were dissolved in 10 mL of chloroform followed by subsequent addition of 0.02 mL of linoleic acid and 0.2 mL of Tween 40. Plant extracts (0.2 mL) were added in the prepared mixture and a control was also run. The resultant mixtures were incubated at 20°C for 15 mins. Chloroform was removed using the rotary evaporator at 40°C and 50 mL of distilled water was added. The obtained mixture was shaken for 2 mins to form emulsion. The absorbance for all samples was measured immediately (0 time) and after incubation of samples for 120 min at 50°C.

$$AA\% = [1 - (A_o - A_t) / (C_o - C_t)] \times 100$$

Where A_o and C_o are the absorbance of sample and control respectively at 0 time. The A_t and C_t are absorbance values of sample and blank after 120 mins of incubation. Positive control contained BHA under same set of conditions.

2.4 Fe-chelating activity

Fe-chelating activity of hydroethanolic leaf extracts was determined following the reported method by Dinis *et al.* (1994). Briefly, 5 mM solution of ferrozine reagent was prepared and 0.2 mL from this were mixed with 100 µL of 2 mM FeSO₄. The methanolic plant extracts at concentration of 2 mg/mL were mixed with reaction mixture followed by an incubation of 10 mins. The absorbance was measured at 562 nm for each sample. Following equation was used to calculate the chelating activity

$$\text{Fe-chelating activity \%} = (A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}} \times 100$$

The control sample contained the all reagents except plant extract, where EDTA was used as positive control.

2.5 Pancreatic lipase inhibition

In-vitro anti-obesity activity of *C. erectus* leaf extracts was performed by measuring the inhibition of pancreatic lipase enzyme. Plant extracts were added to 0.01 M Tris-HCl buffer along with pancreatic lipase. Olive oil was mixed with gum Arabic (dissolved in 0.1 M Tris-HCl buffer having pH 8.0, 20 mM CaCl₂ and 0.5 M NaCl). The lipase inhibition of plant extracts was determined following the previously reported method by Fukumoto *et al.* (1963) with slight modification. To carry out the procedure, 0.5 mL of each plant extract was mixed with 0.2 mL of lipase solution and allowed to stand for 30 minutes at 4°C to allow the reaction to occur completely; following which 2 mL of the substrate was added. The reaction mixture was incubated for 30 mins at 37°C. The reaction was stopped by adding ethanol and acetone in 1:1 composition and the released free fatty acids (FFA) were titrated against 0.02 M NaOH till the pH became 9.4. All the measurements were performed in triplicates and percent inhibition was calculated by the following equation:

$$\% \text{Inhibition} = 100\% - [(V_s / V_c) \times 100]$$

Where, 100% is the enzymatic activity of control. The V_s and V_c represent the volume of base used in titration for sample and control, respectively.

2.6 ¹H-NMR based prediction of metabolites

The most active fraction was subjected to metabolite

prediction. Initially, 25 mg of *C. erectus* most potent leaf extract were mixed with $\text{CH}_3\text{OH}-d_4$ (no internal standard) and with 0.375 mL of KH_2PO_4 buffer (pH=60) in D_2O containing TSP (0.1%). After 1 minute of vortex, ultrasonication was made for 15 mins at 35°C . The solution was subjected to centrifugation at 13000 rpm for 10 mins. Supernatant in amount of 600 μL was collected in NMR tube and $^1\text{H-NMR}$ analysis was performed. INOVA 500 MHz spectrometer (Varian Inc, CA), at 499.887 MHz frequency was used for spectrum generation at 26°C with tetramethylsilane (TMS) was the internal standard. The spectrum was bucketed with MestRenova 11.0 and Chemomx software.

2.7 Statistical analysis

The findings were subjected to analysis of variance with Tukey's test to evaluate the level of significant difference. Minitab 17.0 statistical software was used. The analyses were carried out in triplicate.

3. Results

3.1 Total antioxidant power

The antioxidant potential of plant extracts is represented in Figure 1. High values of absorbance indicated good antioxidant potential of extracts. The 60% ethanolic extract was the most promising one in terms of latent antioxidant capacity having TAP of 223.0 ± 3.27 mg ASE/g PE but less than exhibited by BHA (309.16 ± 3.90 mg ASE/g PE). The aqueous extract exhibited lowest TAP value i.e., 47.33 ± 2.88 mg ASE/g PE among all fractions.




Figure 1. Total antioxidant power of *C. erectus* leaf extracts in comparison with BHA. Different alphabets on the bars are statistically significant at $p < 0.05$.

Statistical analysis indicated significant level of difference ($p < 0.05$) among the TAP values of 60% ethanolic fraction when compared with remaining fractions. The pure ethanolic, 80% and 40% ethanolic fractions were statistically non-significant.

3.2 β -carotene bleaching assay

The comparative evaluation of hydroethanolic leaf extracts of *C. erectus* ranked 60% extract as the most efficient to hinder the bleaching of β -carotene (Figure 2).




Figure 2. Comparative inhibition of β -carotene bleaching by *C. erectus* leaf extracts. Different alphabets on the bars are statistically significant at $p < 0.05$.

The percentage peroxide inhibition for 60% ethanolic extract was $81.39 \pm 2.11\%$ being statistically non-significant from BHA ($89.41 \pm 2.45\%$) which is a strong and efficient synthetic antioxidant ($p < 0.05$). Further comparison elaborated that 60% ethanolic extract was associated with most promising antioxidant attributes among all other solvent fractions. This discriminatory status might be due to variable distribution of phytochemicals in extracts.

3.3 Fe-chelating activity




Figure 3. Fe-chelating activity of *C. erectus* leaf extracts. Different alphabets on the bars are statistically significant at $p < 0.05$.

The results of Fe-chelating activity are presented in Figure 3. The 60% ethanolic extract of *C. erectus* showed maximum metal chelating activity of $68.21 \pm 1.17\%$ in terms of inhibition percentage. The statistical analysis revealed that the metal chelating activity for 60% ethanolic extract was significantly higher than the other respective extracts ($p < 0.05$).

3.4 Pancreatic lipase inhibition

The current investigation regarding PLI presented 60% ethanolic extract as the most efficient anti-obesity fraction followed by 80% ethanolic extract. Statistical analysis indicated the significant level of difference among the percentage inhibition values of extracts, however the PLI for 40% and 80% ethanolic extracts was statistically non-significant ($p < 0.05$). The standard drug orlistat showed maximum PLI with value of 66.65 ± 0.68 and no extract could match the standard drug (Figure 4).

However, PLI of $50.60 \pm 1.47\%$ by 60% ethanolic extract was the highest among extracts and could be considered as promising and effective.




Figure 4. Pancreatic lipase inhibitory action of *C. erectus* leaf extracts and orlistat. Different alphabets on the bars are statistically significant at $p < 0.05$.

3.5 $^1\text{H-NMR}$ based prediction of metabolites




Figure 5. The $^1\text{H-NMR}$ spectrum of 60% ethanolic leaf extract of *C. erectus*.

The $^1\text{H-NMR}$ spectrum of 60% ethanolic extracts of *C. erectus* is represented as Figure 5. The chemical shift values (δ_{H}) corresponded to the nature of metabolites present in leaf extract were highlighted in the spectrum. The $^1\text{H-NMR}$ spectrum is split into aromatic region and aliphatic region. The aromatic region corresponds to the peaks associated with protons of phenolic regime while aliphatic region is used for primary metabolites like sugars, amino acids and fatty acids. The regions ranging

from δ_{H} 6-9 ppm is defined as aromatic region and is used for the identification of polyphenols including flavonoids. The region having δ_{H} 3.5 ppm to 4 ppm and 1-2.5 ppm were categorized as carbohydrate and organic acid regions, respectively. Organic acid region may have peaks for amino acids and fatty acids; however, this region is typically named as amino acid region (Mediani et al., 2012; Mediani et al., 2015).

4. Discussion

Antioxidant activity is a basic and preliminary tool to assess the possible medicinal potential of plants and metabolites. The 60% ethanolic leaf extract of *C. erectus* exhibited highest *in-vitro* antioxidant activities. The β -carotene bleaching based antioxidant potential of 60% ethanolic extract was much higher than recently reported inhibition percentage of 17.88 ± 3.13 for ethanolic extract of *Bromelia laciniosa* (de Oliveira-Júnior et al., 2017). The elevated concentrations of extracts exhibited high antioxidant potential which might be due to higher contents of bioactive constituents mainly phenolic and flavonoids, generally responsible for antioxidant, antiradical and antidiabetic properties of plants (Raza et al., 2013; Evary et al., 2018). Plants are rich sources of natural lipase inhibitors, and hence may provide a workable choice to control obesity along with other associated disorders like diabetes (Jaradat et al., 2017). The lipid metabolism plays a prominent role in energy homeostasis of body. Dietary intake of high fat and high-calorie diets are responsible for weight gain and fat deposition. Lipase enzyme hydrolyzes triglycerides in intestine and facilitates their absorption in body leading to high-fat incorporation. It is very difficult to amend the existing lifestyle and prevailing dietary habits to a great extent for obesity management. On the other hand, complimentary use of synthetic anti-obesity drugs is a matter of concern due to their questionable safety. Under such circumstances, the natural lipase inhibitors are valuable tool for obesity management and plants can serve the purpose smoothly and effectively. The inhibition of pancreatic lipase by plant extracts was probably due to the structural interaction among bioactive metabolites and enzymes. The biologically active substances in extracts may bind with the active sites of pancreatic lipase to modify its functions. Such functional modifications result in reduced enzymatic activity which in turn controls the post-prandial lipidemia (Martinez-Gonzalez et al., 2017).

Iron is required by human body for routine body function but sometimes its excess may be harmful to living system. The ROS production due to iron load may lead to diabetes mellitus making iron load as an important reason behind disease pathogenesis. The exact

mode of action of iron in DMT2 initiation is not fully known but it has been reported that excess iron may damage beta cell of pancreas and also develop insulin insensitivity (Simcox and McClain, 2013). Similarly, the function of liver was also reported to be disturbed under iron load conditions which in turn affected glucose homeostasis and iron extraction (Britton *et al.*, 2018). The Fe-chelating activity by plants extracts indicated that this capacity may be another mechanized reason behind the antidiabetic role of *C. erectus* leaves.

The qualitative prediction of metabolites in 60% ethanolic extract (most potent) was performed by ¹H-NMR spectroscopy. The ¹H-NMR based marking of primary and secondary metabolites has emerged as a novel and effective technique to move into metabolomics. Although, plant metabolomics is a very complicated field especially when plant extract is involved instead of a specific compound or fraction. Besides these tough lines, the ¹H-NMR may be employed to confirm the presence of both primary and secondary plant metabolites (Mediani *et al.*, 2012). The ¹H-NMR technique is unbiased, non-destructive and needs no derivatization. This technique can be used for both qualitative and quantitative analysis of metabolites in crude plant extracts (Mediani *et al.*, 2015). The primary metabolites mainly include carbohydrates and amino acids. Secondary metabolites are mainly, phenolic compounds, flavonoids, tannins and alkaloids. These metabolites produced characteristic peaks or signals due to protonic response to applied magnetic field in ¹H-NMR spectrum and from these signals the nature of metabolites may easily be screened out. The numerous peaks in aromatic region confirmed the presence of secondary metabolites (polyphenols). The secondary metabolites are considered as functional tool to treat chronic ailments including diabetes and obesity.

The antioxidant activity, Fe-chelating activity and pancreatic lipase inhibition potential suggested the *C. erectus* leaves as a novel source of value able natural metabolites which were confirmed by ¹H-NMR response. The *C. erectus* may be exploited to expedite the development in field of nutraceuticals and functional foods.

5. Conclusion

The 60% ethanolic leaf extract of *C. erectus* was proved as the most effective fraction to inhibit bleaching of β-carotene, chelation of iron and inhibition of pancreatic lipase. The ¹H-NMR predicted the presence of various important classes of metabolites including polyphenols and organic acids in leaf extract. The outcomes of this work may be extended further for *in*

vivo trials and toxicity evaluation to move into regime of nutraceuticals and functional foods with antioxidant and antiobesity attributes.

Conflict of Interest

It is declared that authors have no conflict of interest.

References

- Britton, L., Bridle, K., Reiling, J., Santrampurwala, N., Wockner, L., Ching, H., Stuart, K., Subramaniam, V.N., Jeffrey, G., St. Pierre, T., House, M., Gummer, J., Trengove, R., Olynyk, J., Crawford, D. and Adams, L. (2018). Hepatic iron concentration correlates with insulin sensitivity in nonalcoholic fatty liver disease. *Hepatology Communications*, 2 (6), 644-653. <https://doi.org/10.1002/hep4.1190>
- de Oliveira-Júnior, R.G., Ferraz, C.A.A., Souza, G.R., Guimarães, G.R.S.A., de Oliveira, A.P., de Lima-Saraiva, S.R.G., Rolim, L.A., Rolim-Neto, P.J. and da Silva Almeida, J.R.G. (2017). Phytochemical analysis and evaluation of antioxidant and photoprotective activities of extracts from flowers of *Bromelia laciniosa* (Bromeliaceae). *Biotechnology and Biotechnological Equipment*, 31(3), 600-605. <https://doi.org/10.1080/13102818.2017.1288073>
- Dinis, T.C., Madeira, V.M. and Almeida, L.M. (1994). Action of phenolic derivatives (acetaminophen, salicylate, and 5-aminosalicylate) as inhibitors of membrane lipid peroxidation and as peroxy radical scavengers. *Archives of Biochemistry and Biophysics*, 315(1), 161-169. <https://doi.org/10.1006/abbi.1994.1485>
- Evary, Y.M., Nugroho, A.E. and Pramono, S. (2018). Comparative study on DPPH free radical scavenging and alpha-glucosidase inhibitory activities of ethanolic extracts from different parts of durian plant (*Durio zibethinus* Murr.). *Food Research*, 3(5), 463 – 468. [https://doi.org/10.26656/fr.2017.3\(5\).085](https://doi.org/10.26656/fr.2017.3(5).085)
- Fukumoto, J., Tsujisaka, Y. and Iwai, M. (1963). STUDIES ON LIPASE. *The Journal of General and Applied Microbiology*, 9(3), 353-361. [https://doi.org/10.26656/fr.2017.3\(5\).085](https://doi.org/10.26656/fr.2017.3(5).085)
- Jaradat, N., Zaid, A., Hussein, F., Zaqzouq, M., Aljammal, H. and Ayeshe, O. (2017). Anti-lipase potential of the organic and aqueous extracts of ten traditional edible and medicinal plants in Palestine; a comparison study with orlistat. *Medicines*, 4(4), 89. <https://doi.org/10.3390/medicines4040089>
- Madhujith, T. and Shahidi, F. (2006). Optimization of the extraction of antioxidative constituents of six barley cultivars and their antioxidant properties. *Journal of Agricultural and Food Chemistry*, 54(21),

- 8048-8057. <https://doi.org/10.1021/jf061558e>
- Martinez-Gonzalez, A.I., Alvarez-Parrilla, E., Díaz-Sánchez, Á.G., de la Rosa, L.A., Núñez-Gastélum, J.A., Vazquez-Flores, A.A. and Gonzalez-Aguilar, G.A. (2017). In vitro inhibition of pancreatic lipase by polyphenols: A kinetic, fluorescence spectroscopy and molecular docking study. *Food Technology and Biotechnology*, 55(4), 519-530. <https://doi.org/10.17113/ftb.55.04.17.5138>
- Mediani, A., Abas, F., Khatib, A., Maulidiani, H., Shaari, K., Choi, Y.H. and Lajis, N. (2012). ¹H-NMR-based metabolomics approach to understanding the drying effects on the phytochemicals in *Cosmos caudatus*. *Food Research International*, 49(2), 763-770. <https://doi.org/10.1016/j.foodres.2012.09.022>
- Mediani, A., Abas, F., Khatib, A., Tan, C.P., Ismail, I.S., Shaari, K., Ismail, A. and Lajis, N.H. (2015). Phytochemical and biological features of *Phyllanthus niruri* and *Phyllanthus urinaria* harvested at different growth stages revealed by ¹H NMR-based metabolomics. *Industrial Crops and Products*, 77, 602-613. <https://doi.org/10.3390/molecules22060902>
- Nascimento, D.K., Souza, I.A., OLiveira, A.F.D., Barbosa, M.O., Santana, M.A., Pereira Junior, D.F., Lira, E.C. and Vieira, J.R. (2016). Phytochemical screening and acute toxicity of aqueous extract of leaves of *Conocarpus erectus* Linnaeus in swiss albino mice. *Anais da Academia Brasileira de Ciências*, 88(3), 1431-1437. <https://doi.org/10.1590/0001-3765201620150391>
- Piyush, G., Mendhe, B., Namrata, S., Milind, P., Anupam, P. and Sahadev, Y. (2010). Antioxidant activity of flavonoidal content of *Butea monosperma*. *International Journal of Pharmaceutical Sciences and Research (IJPSR)*, 1 (4), 85-88. [http://dx.doi.org/10.13040/IJPSR.0975-8232.1\(4\).85-88](http://dx.doi.org/10.13040/IJPSR.0975-8232.1(4).85-88)
- Prieto, P., Pineda, M. and Aguilar, M. (1999). Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E. *Analytical Biochemistry*, 269(2), 337-341. <https://doi.org/10.1006/abio.1999.4019>
- Raza, S.A., Chaudhary, A.R., Mumtaz, M.W., Ghaffar, A., Adnan, A. and Waheed, A. (2018). Antihyperglycemic effect of *Conocarpus erectus* leaf extract in alloxan-induced diabetic mice. *Pakistan Journal of Pharmaceutical Sciences*, 31(2 Suppl), 637-642.
- Raza, S.A., Rashid, A., William, J., Arshed, S.F. and Arshad, M. (2013). Comparison of antioxidant activity of some medicinally important plants from Pakistan. *Acta Scientiarum Polonorum Technologia Alimentaria*, 12(4), 403-410.
- Saleh, E.-S., Bazaid, S., Shohayeb, M., El-Sayed, M. and El-Wakil, E. (2012). Phytochemical studies and evaluation of antioxidant, anticancer and antimicrobial properties of *Conocarpus erectus* L. growing in Taif, Saudi Arabia. *European Journal of Medicinal Plants*, 2(2), 93-112. <https://doi.org/10.1155/2013/345465>
- Saleh, E.-S., Bazaid, S.A. and Sbara, A.-N. (2013). Protective effect of *conocarpus erectus* extracts on CCl₄-induced chronic liver injury in mice. *Global Journal of Pharmacology*, 7, 52-60.
- Shaaban, M.A., Dawod, A.E.A. and Nasr, M.A. (2016). Role of iron in diabetes mellitus and its complications. *Menoufia Medical Journal*, 29(1), 11-16. <https://doi.org/10.4103/1110-2098.178938>
- Shon, M.Y., Kim, T.H. and Sung, N.J. (2003). Antioxidants and free radical scavenging activity of *Phellinus baumii* (Phellinus of Hymenochaetaceae) extracts. *Food Chemistry*, 82(4), 593-597. [https://doi.org/10.1016/S0308-8146\(03\)00015-3](https://doi.org/10.1016/S0308-8146(03)00015-3)
- Simcox, J.A. and McClain, D.A. (2013). Iron and diabetes risk. *Cell Metabolism*, 17(3), 329-341. <https://doi.org/10.1016/j.cmet.2013.02.007>
- Wang, J., Zhang, Q., Zhang, Z., Song, H. and Li, P. (2010). Potential antioxidant and anticoagulant capacity of low molecular weight fucoidan fractions extracted from *Laminaria japonica*. *International Journal of Biological Macromolecules*, 46(1), 6-12. <https://doi.org/10.1016/j.ijbiomac.2009.10.015>
- William, J., John, P., Mumtaz, M. W., Ch, A. R., Adnan, A., Mukhtar, H., Sharif, S. and Raza, S.A. (2018). Antioxidant activity, Hypoglycemic potential and metabolite profiling of *Hyophorbe indica* leaf extract. *Pakistan Journal of Pharmaceutical Sciences*, 31(6 Suppl.), 2737-2742.