

Primary and secondary antioxidant activities of nine edible mushrooms species¹Muhammad Ezzudin, R., ^{1*}Rabeta, M.S., ²Shaida Fariza, S., ²Leong Ooi, K. and ²Nurul Shafiqah, H.*Food Technology Division, School of Industrial Technology, Universiti Sains Malaysia, 11800 Minden, Penang, Malaysia.**²School of Biological Sciences, Universiti Sains Malaysia, 11800 Minden, Penang, Malaysia***Article history:**

Received: 31 March 2018

Received in revised form: 12 June 2018

Accepted: 20 June 2018

Available Online: 22 June 2018

Keywords:Antioxidant activity,
Total phenolic content,
Mushrooms**DOI:**[https://doi.org/10.26656/fr.2017.3\(1\).077](https://doi.org/10.26656/fr.2017.3(1).077)**Abstract**

In this study, nine different edible mushrooms, namely bunapi, maitake, bunashimeji, abalone, oyster, button, white, enoki and kukur evaluated for antioxidant activities and total phenolic content (TPC). Specifically, three different antioxidant assays which were 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radical scavenging, ferric reducing antioxidant power (FRAP) and metal chelating (MC) used. From the evaluation, button mushrooms were identified to scavenge free radicals and reduce lipid peroxidation with the lowest EC₅₀ of 23.40±2.56 mg/mL in methanol extract. Besides, the highest ferric reducing antioxidant power and metal chelating activities were exhibited by button mushroom in water extract at 25.47±2.52 mg/mL and enoki in methanol extract at 1.00±0.16 mg/mL, respectively. On the other hand, the highest contents of total phenolics were recorded in a water extract of kukur with 211.35±11.89 µg GAE/g extract and methanol extract of maitake with 400.28±11.35 µg GAE/g extract. However, there was no correlation between the antioxidant activity and total phenolic content because the activity of DPPH, FRAP and metal chelating revealed different lowest EC₅₀ value when associated with different species of mushrooms. Hence, this study showed the antioxidant values of nine edible mushrooms species, which can serve as a useful database for the selection of mushrooms in food, pharmaceuticals and nutraceuticals industries.

1. Introduction

In general terms, primary antioxidant or chain breaking antioxidant is a compound that reacts with radicals to convert them into more stable products. For instance, alkyl peroxy radical, ROO• is a significant lipid radical at normal oxygen pressures that functions as an oxidizing agent which may be directly converted into a hydroperoxide by a hydrogen donor (Gordon, 1990). Conversely, secondary antioxidants are compounds that inhibit the autoxidation of lipids and interrupt the autoxidation chain by converting free radicals into more stable species (Lobo *et al.*, 2010). Among the mechanisms available which could inhibit the rate of autoxidation are metal ion binding, oxygen scavenging, UV radiation absorption or singlet oxygen deactivation and decomposition of hydroperoxides into non-radical species (Lü *et al.*, 2010). In fact, metal ions and reducing agents such as ascorbic acid have identified as active sequestering agents in revealing secondary antioxidant in the presence of tocopherols or other phenolic compounds (Gordon, 1990).

Mainly, a mushroom is a macrofungus with a distinctive fruiting body which can be either epigeous (above ground) or hypogeous (underground) and visible to our naked eyes (Miles and Chang, 2004). Mushrooms can be classified into six different groups according to their morphological characteristics (Fitts, 1961). To this classification, the groups are basidiomycetes (spore or reproductive bodies are naked), ascomycetes (spores enclosed in sacs or asci), physcomycetes, myxomycetes (slime moulds), saccharomycetes (yeast fungi) and schizomycetes. Fungi can live in three different conditions which are saprophytic, parasitic and symbiotic. In this context, saprophytic mushrooms will use their enzymes to break down the organic matter inside mushrooms. For instance, one of the examples of saprophytic fungi is the shaggy ink cup (*Coprinus comatus*). Secondly, parasitic mushrooms are mushrooms that grow on trees and absorb nutrients from the trees. One species which falls this category is The Honey Fungus (*Armillaria mellea*). Thirdly, symbiotic fungi mushrooms are essential components of stream ecosystems and critical intermediaries in the transfer of carbon, nutrients, and energy to higher trophic levels

*Corresponding author.

Email: rabeta@usm.my

(Gulis and Bärlocher, 2017).

Furthermore, the antioxidant activity of different edible mushroom extracts plays a significant role in absorbing and neutralizing free radicals in a deleterious molecule, decomposing the peroxide and quenching singlet and triplet oxygen (Osawa, 1994). In fact, based on a proximate analysis conducted by Wang *et al.* (2014), edible mushrooms were discovered to be rich in carbohydrate, protein and essential minerals (potassium, calcium, and magnesium). Besides, ascorbic acid, tocopherols, phenolics, flavonoids, and carotenoids are among the group of compounds present in mushrooms with antioxidant activity (Ferreira *et al.*, 2009). Moreover, mushrooms have acknowledged as a nutritious and healthy food due to the bioactive compounds namely phenolic which have the ability in scavenging free radical, reducing ferric to ferrous ion and chelating metal ion (Decker, 1997). Additionally, edible mushrooms could be used directly to increase our dietary antioxidant intake (Kozarski *et al.*, 2015). Enokitake (*Flammulina velutipes*), shiitake (*Lentinula edodes*) and split gill mushrooms (*Schizophyllum commune*) were locally cultivated in Malaysia while button mushrooms (*Agaricus bisporus*), Nameko (*Pholiota nameko*) and *Hypsizyguis* sp. were imported from other countries (Lee *et al.*, 2009).

To date, studies to determine the antioxidant activities in mushrooms remain insufficient. Studies on the antioxidant activity of mushrooms in Malaysia are mostly focused on primary antioxidant (Arbaayah and Kalsom, 2013) and less on secondary antioxidant activity. Moreover, samples of mushrooms for research were limited and only the well-known species such as *Pleurotus* spp. and shiitake (*Lentinula edodes*) were given preference. Therefore, this comparative study was performed to determine the primary and secondary antioxidant activities of nine edible mushrooms species commonly consumed in Malaysia and identify mushrooms with the highest antioxidant activity.

2. Materials and methods

2.1 Samples preparation

Nine species of edible mushrooms were bought from three different supermarkets around Penang Island which are Tesco Sungai Dua, Queensbay Mall and Giant Hypermarket in Bukit Gedung. Blanched samples were blended, and the juices were squeezed out, prior to analysis.

2.2 Water extraction

The extraction method was prepared according to the procedure used by Ooi *et al.* (2011) with slightly

modification. The sample was ground using a Smart Lady blender juice and was then put on the muslin cloth. The juice was squeezed out and was put in 250 mL beaker. The juice was then put into a 50 mL centrifuge tubes and were kept in the -20°C laboratory freezer prior to analysis. The residue of samples was extracted with methanol. All extractions were performed in triplicate.

2.3 Methanol extraction

The residue was weighed using electronic beam balance (BEL Engineering) and put it into 250 mL beaker. Methanol (100%) was added to the beaker 5 times higher than the weight of the filtrate. The ratio of the filtrate to the 100% methanol added was 1:5. Then, the mixture was boiled for about 20 minutes. After that, the extract was filtered using muslin cloth in 250 mL beaker. Finally, the methanol extract was added to a 50 mL centrifuge tube prior to analysis (Ooi *et al.*, 2011). All extractions were carried out in triplicate.

2.4 Determination of DPPH assay

About 50 µL of the prepared extracts were loaded to each well of the 96-microwell plate. A 150 µL of 0.3 mM 1,1-diphenyl-2-picrylhydrazyl was pipetted into each well that contained the extracts using multi-channel pipette in the dark room. The total amount of solution in each well was 200 µL. The quercetin was used as a positive control while dimethyl sulfoxide (DMSO) as a negative control. Then, 96-microwell plate was wrapped in aluminium foil before incubated for 30 minutes at 37°C. The absorbance reading was taken after 30 minutes using Multiskan Spectrum (Thermo Scientific) at wavelength 515 nm (Brand-Williams *et al.*, 1995). The results obtained was compared with the positive control, quercetin. The purple colour of the DPPH will turn to yellow when the sample had an antioxidant activity. The ability of the extracts to scavenge at 50% of the DPPH, EC₅₀ was determined from the graph plotted in GraphPad Prism software. All experiments were carried out in triplicate.

2.5 Determination of FRAP assay

A 50 µL of the extracts was pipetted into the plates. Then, 150 µL of FRAP reagent was added directly to each well Trolox was used as a positive control while DMSO as a negative control of the experiment. Then, 96 -microwell plate was incubated at 37°C for 30 minutes before measuring the absorbance. After 30 mins incubated in the incubator, the absorbance of the extracts was measured using Multiskan Spectrum (Thermo Scientific) at a wavelength of 593 nm (Benzie and Strain, 1996). The absorbance of the extracts was compared with the positive control, Trolox. The colour of the FRAP turns to deep blue when there was an antioxidant

activity. The EC₅₀ was determined from the graph that was plotted in GraphPad Prism software. All experiments were performed in triplicate.

2.6 Determination of metal chelating assay

Approximately 50 µL of each extract was incubated with 5 µL ferrous chlorides and 130 µL of deionized water for 10 mins at 37°C. The mixture was added with 15 µL of ferrozine which initiated the reactions. The mixture then was incubated for 10 mins at 37°C. After 10 mins of incubation, the absorbance of the extracts was quickly read by Multiskan Spectrum (Thermo Scientific) at wavelength 562 nm (Dinis *et al.*, 1994). The absorbance of the extracts was compared to the Ethylenediaminetetraacetic acid (EDTA) as a positive control. DMSO was used as the negative control and ferrous chloride was used as background in this test. The colour of the metal chelating assay will turn from bright purple to pale purple when there was an antioxidant activity. The ability of the ferrous ion to be 50% chelated by the extracts was determined as EC₅₀ from the graph plotted in GraphPad Prism software. All experiments were carried out at least three replicates.

2.7 Determination of total phenolic content

Approximately 50 µL of prepared extracts was loaded to the 96-microwell plate. A 25 µL of the Folin-Ciocalteu (FC) solution was added to the extracts (Singleton *et al.*, 1999). The mixture was incubated for 5 mins at room temperature (28°C). After 5 minutes of incubation, 25 µL of 2% sodium carbonate (NaCO₃) solution was added to the mixture followed by 100 µL of distilled water to a final volume of 200 µL per well. Then, the mixture was incubated for about 30 mins at room temperature (28°C) before the reading was taken using Multiskan Spectrum (Thermo Scientific) at wavelength of 750 nm. The absorbance obtained was compared to the positive control, gallic acid. All experiments were performed in triplicate. The results expressed as microgram gallic acid equivalent per gram extracts (µg GAE /g extract).

2.8 Statistical analysis

The results expressed as the mean standard deviation (SD). Statistical analysis of data was carried out using SPSS version 20.0. Significant differences between means calculated by using Duncan's multiple range tests at a level of $p < 0.05$.

3. Results and discussion

3.1 DPPH scavenging activity of the samples

As shown in Table 1, for the DPPH assay, white mushrooms (97.05±19.21%) and bunashimeji (92.14±2.86%) possessed the highest percentage of scavenging activity in aqueous extract followed by bunapi, oyster, button, kukur, abalone, and enoki. The lowest scavenging activity was recorded by maitake with the percentage value at 13.39±1.96%.

The methanolic button mushroom extract exhibited the highest scavenging free radical activity with the value of 77.62±8.11%. Next, enoki was ranked second highest at 70.70±3.58%. In this study, bunapi, bunashimeji, oyster, maitake and abalone mushrooms were identified to possess moderate percentage values ranging from 37.39±3.32% to 24.67±1.53%. Also, white mushroom and kukur had the least effective scavenging activity with the lowest percentage of 17.81±1.15% and 15.09±1.34%, respectively.

In contrast, button mushroom was exhibited the lowest EC₅₀ value of 23.40±2.56 mg/mL compared to other mushrooms both in aqueous and methanolic extracts. However, the comparison of methanol and water extracts did not show any significant difference ($p > 0.05$). The sequence of antioxidant activity in DPPH test of aqueous and methanolic extracts was as follows:

Button (methanol) > Enoki (methanol) > Bunapi (water) > Bunashimeji (water) > White (water)

In this study, the result agrees with a study done by Fu *et al.* (2002), where the best result of scavenging

Table 1. The EC₅₀ values of the DPPH scavenging activity of water and methanol extracts.

Samples	Water Extract		Methanol Extract	
	Percentage of scavenging activity (%)	EC ₅₀ of DPPH Scavenging activity (mg/mL)	Percentage of scavenging activity (%)	EC ₅₀ of DPPH Scavenging activity (mg/mL)
Bunapi	69.18±3.73	68.31±3.55	37.39±3.32	*ND
Maitake	13.39±1.96	*ND	25.16±2.33	*ND
Bunashimeji	92.14±2.86	133.10±34.43	34.97±0.51	*ND
Abalone	37.19±3.86	*ND	24.67±1.53	*ND
Oyster	48.63±2.34	*ND	31.50±1.76	*ND
Button	38.59±22.05	*ND	77.62±8.11	23.40±2.56
White	97.05±19.21	165.97±15.25	17.81±1.15	*ND
Enoki	34.82±3.73	*ND	70.70±3.58	35.86±1.58
Kukur	38.01±0.43	*ND	15.09±1.34	*ND
Quercetin	78.15±0.70	-	78.15±0.70	-

Key: *ND = Not Determined

Table 2. The EC₅₀ values of the ferric reducing activity of water and methanol extracts.

Samples	Water Extract		Methanol Extract	
	Percentage of reducing activity (%)	EC ₅₀ of FRAP reducing activity (mg/mL)	Percentage of reducing activity (%)	EC ₅₀ of FRAP reducing activity (mg/mL)
Bunapi	*ND	15.95±1.06	*ND	16.54±1.24
Maitake	*ND	39.46±0.34	*ND	14.63±2.52
Bunashimeji	*ND	14.99±0.89	*ND	16.39±0.98
Abalone	*ND	16.35±0.10	*ND	11.19±1.36
Oyster	*ND	24.42±6.63	*ND	15.16±1.50
Button	25.47±2.52	84.60±0.00	*ND	43.91±0.49
White	*ND	17.06±0.08	*ND	20.35±0.25
Enoki	*ND	12.86±0.11	*ND	24.43±1.24
Kukur	*ND	48.69±2.52	*ND	12.57±1.24
Trolox	73.25±0.34	-	73.25±0.34	-

Key: *ND = Not Determined

activity observed in button mushrooms, *Agaricus bisporus* when methanol extract used. This result might be due to the significant amount of phenolic amino acids such as tyrosine, L-glutaminy-4-hydroxybenzene and ascorbic acid which promoted the scavenging activity (Choi and Sapers, 1994).

3.2 FRAP assay of the samples

In Table 2, button possessed the highest reducing activity with the value of 84.60% at the highest concentration in water extract then followed by kukur and maitake at 48.69±2.52% and 39.46±0.34%, respectively. Next, oyster produced a percentage of 24.42±6.63% when tested at the highest concentration. On the other hand, the average percentage of reducing activity was white, abalone, bunapi, and bunashimeji at 17.06±0.08%, 16.35±0.10%, 15.95±1.06% and 14.99±0.89%, respectively. In fact, the lowest percentage of ferric reducing activity was exhibited by enoki only at 12.86±0.11%.

Button mushroom also exhibited the highest ferric reducing activity when examined with methanolic extract which resulted in the value of 43.91±0.49%. The list then followed by enoki and white mushrooms with the percentage of 24.43±1.24% and 20.35±0.251%, respectively. Next, bunapi, bunashimeji, oyster and

maitake possessed the moderate percentage values of 16.54±1.24%, 16.39±0.98%, 15.16±1.50% and 14.63±2.52%, respectively. For the values obtained, kukur and abalone classified as mushrooms with the least ferric reducing activity with the lowest percentage at 12.57±1.24% and 11.19±1.36%, respectively.

In this assay, the highest antioxidant potential with the lowest EC₅₀ value also exhibited by button mushroom. However, it should note that it was only in the water extract that activity occurred with the EC₅₀ value of 25.47±2.52%. Conversely, the EC₅₀ for the other mushrooms extracts were unable to be determined. For instance, at ($p < 0.05$), in relation to this study, there was a significant difference between water and methanol extracts EC₅₀. According to Gan *et al.* (2013), the water extract of button mushroom is an effective source of activity reduction compared to ethanol. In this context, this might be due to the presence of extraction solvent and test system used (Fu *et al.*, 2010).

3.3 Metal chelating assay of the samples

As demonstrated in Table 3, white mushrooms exhibited the highest percentage of metal chelating activity with the value of 99.95±0.50% when tested at the highest concentration in water extract. The kukur mushroom followed this with a reading of 99.27±2.79%

Table 3. The EC₅₀ values of the metal chelating activity of water and methanol extracts.

Samples	Water Extract		Methanol Extract	
	Percentage of chelating activity (%)	EC ₅₀ of metal chelating activity (mg/mL)	Percentage of chelating activity (%)	EC ₅₀ of metal chelating activity (mg/mL)
Bunapi	95.00±4.84	3.34±0.22	90.98±4.82	1.06±0.13
Maitake	82.33±6.65	7.54±0.31	98.45±0.57	2.89±0.06
Bunashimeji	88.23±6.52	3.82±0.15	99.45±0.27	1.15±0.12
Abalone	94.48±0.18	50.30±0.74	99.60±4.87	1.95±0.29
Oyster	91.12±2.30	6.38±0.59	92.68±0.25	5.11±0.17
Button	89.51±0.61	30.34±0.57	94.42±9.03	1.26±0.06
White	99.95±0.5	2.42±0.16	99.23±10.04	1.51±0.08
Enoki	98.38±0.20	30.62±0.75	99.49±0.79	1.00±0.16
Kukur	99.27±2.79	25.54±2.04	35.64±0.18	*ND
EDTA	95.86±0.18	-	95.86±0.18	-

Key: *ND = Not Determined

at the highest concentration. Meanwhile, enoki, bunapi, abalone, oyster, button and bunashimeji mushrooms produced a relatively average percentage of metal chelating activity. Concerning this finding, maitake was identified to have the lowest metal chelating activity with the percentage value of $82.33 \pm 6.65\%$.

On the other hand, in methanol extract abalone mushroom possessed the highest metal chelating activity with the value of $99.60 \pm 4.87\%$ at the highest concentration. Then, the order continued with enoki and bunashimeji at $99.49 \pm 0.79\%$ and $99.45 \pm 0.27\%$, respectively. Lastly, the white mushrooms recorded a percentage of $99.23 \pm 10.04\%$ at its highest concentration. Moreover, it also discovered that mushrooms with the average percentage of metal chelating activity were enoki, maitake, button, oyster, and bunapi ranging from $99.23 \pm 10.04\%$ to $90.98 \pm 4.83\%$, respectively. In connection with this study, the lowest percentage of metal chelating activity was exhibited by kukur only at $35.64 \pm 0.18\%$.

Additionally, enoki mushroom (*Flammulina velutipes*) of methanol extract produced the lowest EC_{50} value of 1.00 ± 0.16 mg/mL. In this respect, this reading might potentially influence by the presence of solid active compounds presence such as phenol and flavonoid that exhibit favorable chelating of Fe^{2+} activity (Ebrahimzadeh et al., 2008). Particularly, bunapi, bunashimeji and white mushrooms of methanol extract showed lower EC_{50} value with 1.06 ± 0.13 mg/mL, 1.15 ± 0.12 mg/mL and 1.51 ± 0.08 mg/mL, respectively compared to water extract at 3.34 ± 0.22 mg/mL, 3.82 ± 0.15 mg/mL and 3.66 ± 0.12 mg/mL, respectively. Based on the results recorded, there were no significant differences found between water and methanol extracts of EC_{50} at ($p > 0.05$).

3.4 Total phenolic content

Table 4 showed that among the water extracts, the highest total phenolic content exhibited by kukur and this was followed by maitake which produced readings of 211.35 ± 11.89 μ g GAE/g extract and 129.44 ± 1.06 μ g GAE/g extract, respectively. Total phenolics are the major naturally occurring antioxidant components found in Kukur followed by Maitake in water extract. About this study, the lowest total phenolics content was enoki mushrooms with 30.97 ± 1.78 μ g GAE/g extract while the other extracts showed moderate of total phenolic contents. Moreover, among the methanol extracts, the highest total phenolic content was exhibited by the maitake mushrooms at 400.28 ± 11.35 μ g GAE/g extract, and the lowest was kukur at 3.58 ± 0.70 μ g GAE/g extract. On the other hand, the remaining showed moderate of total phenolic content.

Table 4. Total phenolic content of water and methanol extracts.

Samples	Total phenolic content (μ g GAE/g extract)	
	Water Extract	Methanol Extract
Bunapi	84.97 ± 0.61	36.34 ± 3.64
Maitake	129.44 ± 1.06	400.28 ± 11.35
Bunashimeji	69.52 ± 0.53	74.69 ± 0.66
Abalone	45.77 ± 0.22	15.16 ± 1.02
Oyster	38.56 ± 2.90	15.42 ± 1.54
Button	33.13 ± 0.49	56.26 ± 1.50
White	74.69 ± 0.66	16.18 ± 0.73
Enoki	30.97 ± 1.78	35.49 ± 0.45
Kukur	211.35 ± 11.89	3.58 ± 0.70

Key: *ND = Not Determined

In this study, the highest total phenolic content was displayed by methanol extract of Maitake mushrooms (*Grifola frondosa*) at 400.28 ± 11.35 μ g GAE/g extract. Besides, most mushrooms of the water extract also exhibited higher total phenolic contents compared to methanol extracts such as kukur and bunapi at 211.35 ± 11.89 μ g GAE/g extract and 84.97 ± 0.61 μ g GAE/g extract, respectively. In this respect, it deduced that there is a significant difference between water and methanol extracts EC_{50} at ($p < 0.05$).

Contrariwise, a study conducted by Mau et al. (2002) had shown that methanolic extract of maitake mushrooms contains moderate content of total phenolics compared to the others three mushrooms which are *Dictophora indusiata*, *Hericium erinaceus*, and *Tricholoma giganteum*. However, these mushrooms not used in this study. In other words, the differences in total phenolic content suggest that they might have contributed by the sample processing method and extraction method or the cultivar studied. A group of researchers also highlighted that the major phenolic compounds found in mushrooms are phenolic acids (Ferreira et al., 2009). The nine edible mushrooms used in this study could be categorized into three groups based on their phenolics content as low, moderate, and high phenolic species (Puttaraju et al., 2006). Thus, it can be concluded that the total phenolic content of the methanol extract of maitake was the highest compared to the other mushrooms in water extract.

4. Conclusion

In conclusion, it can be deduced from the results that there is no correlation between the antioxidant activity and total phenolic content because the activity of DPPH, FRAP and metal chelating showed that the different mushrooms exhibited the lowest EC_{50} value with phenolics content. Therefore, the 50% (v/v) water extract of *white mushroom* can be used as a potential natural antioxidative source in food industries. Based on the

results of this study, a few suggestions were established to improve this research in the future. For instance, one of the recommendations includes increasing the number of mushrooms samples by including various species such as Shiitake (*Lentinula edodes*), Jeli Hitam (*Auricularia* sp.) and Ganoderma (*Ganoderma* sp.) to increase the validity of the results regarding the antioxidant activity. Hence, this study revealed the antioxidant values of nine edible mushrooms species, which can serve as a useful database for the selection of mushrooms in food, pharmaceuticals and nutraceuticals industries.

Conflict of interest

The authors declared no conflict of interest.

Acknowledgment

School of Biological Sciences, Universiti Sains Malaysia 11800 Minden, Penang, Malaysia contributed to the funding of this research.

References

- Arbaayah, H. and Kalsom, Y. (2013). Antioxidant properties in the oyster mushrooms (*Pleurotus* spp.) and split gill mushroom (*Schizophyllum commune*) ethanolic extracts. *Mycosphere*, 4(4), 661-673.
- Benzie, I.F. and Strain, J.J. (1996). The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Analytical Biochemistry*, 239(1), 70-76. <https://doi.org/10.1006/abio.1996.0292>
- Brand-Williams, W., Cuvelier, M.-E. and Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *LWT-Food Science and Technology*, 28(1), 25–30. [https://doi.org/10.1016/S0023-6438\(95\)80008-5](https://doi.org/10.1016/S0023-6438(95)80008-5)
- Choi, S.W. and Sapers, G.M. (1994). Effects of washing on polyphenols and polyphenol oxidase in commercial mushroom (*Agaricus bisporus*). *Journal of Agriculture and Food Chemistry*, 42(10), 2286-2290. <https://doi.org/10.1021/jf00046a038>
- Decker, A. (1997). Phenolics: prooxidants or antioxidants? *Nutrition Review*, 55(11), 396-398.
- Dinis, T.C., Madeira, V.M. and Almeida, L.M. (1994). Action of phenolic derivatives (acetaminophen, salicylate, and 5-aminosalicylate) as inhibitors of membrane lipid peroxidation and as peroxy radical scavengers. *Archives of Biochemistry and Biophysics*, 315(1), 161-169. <https://doi.org/10.1006/abbi.1994.1485>
- Ebrahimzadeh, M.A., Pourmorad, F. and Bekhradnia, A.R. (2008). Iron chelating activity, phenol and flavonoid content of some medicinal plants from Iran. *African Journal of Biotechnology*, 7(18), 3188-3192
- Ferreira, I.C., Barros, L. and Abreu, R. (2009). Antioxidants in wild mushrooms. *Current Medicinal Chemistry*, 16(12), 1543-1560. <https://doi.org/10.2174/092986709787909587>
- Fu, H.Y., Shieh, D.E. and Ho, C.T. (2002). Antioxidant and free radical scavenging activities of edible mushrooms. *Journal of Food Lipid*, 9(1), 35-43. <https://doi.org/10.1111/j.1745-4522.2002.tb00206.x>
- Fitts, D.D. (1961). The mushrooms: edible and otherwise its habitat and its time of growth. 12th ed. New York: Hafner Publishing Company.
- Gan, C., Nurul Amira, B. and Asmah, R. (2013). Antioxidant analysis of different types of edible mushrooms (*Agaricus bisporous* and *Agaricus brasiliensis*). *International Food Research Journal*, 20(3), 1095-1102.
- Gordon, M.H. (1990). The mechanism of antioxidant action *in Vitro*. In Hudson, B.J.F. (Ed.) Food Antioxidants. Elsevier Applied Food Science Series, p. 1-18. Dordrecht: Springer. https://doi.org/10.1007/978-94-009-0753-9_1
- Gulis, V. and Bärlocher, F. (2017). Fungi: Biomass, Production and Community Structure. In Hauer, R.F. and Lamberti, G.A. (Eds.) Methods in Stream Ecology. Vol. 1, 3rd ed., p. 177-192. USA: Academic Press. <https://doi.org/10.1016/B978-0-12-416558-8.00010-X>
- Kozarski, M., Klaus, A., Jakovljevic, D., Todorovic, N., Vunduk, J., Petrović, P., Niksic, M., Vrvic, M.M. and van Griensven, L. (2015). Antioxidants of Edible Mushrooms. *Molecules*, 20(10), 19489-19525. <https://doi.org/10.3390/molecules201019489>
- Lee, S., Chang, Y. and Noraswati, M. (2009). Utilization of macrofungi by some indigenous communities for food and medicine in Peninsular Malaysia. *Forest Ecology and Management*, 257(10), 2062-2065. <https://doi.org/10.1016/j.foreco.2008.09.044>
- Lobo, V., Patil, A., Phatak, A. and Chandra, N. (2010). Free radicals, antioxidants and functional foods : Impact on human health. *Pharmacognosy Review*, 4 (8), 118–126. <https://doi.org/10.4103/0973-7847.70902>
- Lü, J.-M., Lin, P.H., Yao, Q. and Chen, C. (2010). Chemical and molecular mechanisms of antioxidants: experimental approaches and model systems. *Journal of Cellular and Molecular Medicine*, 14(4), 840–860. <https://doi.org/10.1111/j.1582-4934.2009.00897.x>
- Mau, J. L., Lin, H.C. and Song, S.F. (2002). Antioxidant

- properties of several specialty mushrooms. *Food Research International*, 35(6), 519-526. [https://doi.org/10.1016/S0963-9969\(01\)00150-8](https://doi.org/10.1016/S0963-9969(01)00150-8)
- Miles, P.G. and Chang, S.T. (2004). Mushrooms: cultivation, nutritional value, medicinal effect, and environmental impact. USA: CRC Press.
- Ooi, K.L., Muhammad, T.S.T., Tan, M.L. and Sulaiman, S.F. (2011). Cytotoxic, apoptotic and anti- α -glucosidase activities of 3, 4-di-O-caffeoylquinic acid, an antioxidant isolated from the polyphenolic-rich extract of *Elephantopus mollis* Kunth. *Journal of Ethnopharmacology*, 135(3), 685-695. <https://doi.org/10.1016/j.jep.2011.04.001>
- Osawa, T. (1994). Novel natural antioxidant for utilization in food and biological systems. *Postharvest Biochemistry of Plant Food Materials in Tropics*, p. 241-251. Tokyo: Japan Scientific Press
- Puttaraju, N.G., Venkateshaiah, S.U., Dharmesh, S.M., Nanjaraj Urs, S.M. and Somasundaram, R. (2006). Antioxidant Activity of Indigenous Edible Mushrooms. *Journal of Agricultural and Food Chemistry*, 54(26), 9764-9772. <https://doi.org/10.1021/jf0615707>
- Singleton, V.L., Orthofer, R. and Lamuela-Raventós, R. M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*, 299, 152-178. [https://doi.org/10.1016/S0076-6879\(99\)99017-1](https://doi.org/10.1016/S0076-6879(99)99017-1)
- Wang, X.M., Zhang, J., Wu, L.H., Zhao, Y.L., Li, T., Li, J.Q., Wang, Y.Z. and Liu, H.-G. (2014). A mini-review of chemical composition and nutritional value of edible wild-grown mushroom from China. *Food Chemistry*, 151, 279-285. <https://doi.org/10.1016/j.foodchem.2013.11.062>